




New Mexico State University
All About Discovery!
nmsu.edu

COHERENT at the Spallation Neutron Source

Robert L. Cooper
New Mexico State University
on behalf of the COHERENT
Collaboration



Outline

- Physics Motivation for Coherent Elastic Neutrino-Nucleus Scattering (CE ν NS)
- How to measure CE ν NS
 - Production of neutrinos at SNS
 - Detection with multiple technologies
 - Suppression of background neutrons
- Status of COHERENT and other activities



"Wait a minute! Isn't anyone here a real sheep?"

CE ν NS (pronounced sevens)

- Coherent Elastic Neutrino-Nucleus Scattering
- Flavor blind Neutral Current process that scatters the entire nucleus as a whole
- To probe a “large” nucleus (few $\times 10^{-15}$ m)

$$E_\nu \lesssim \frac{hc}{R_N} \cong 50 \text{ MeV}$$

- Recoiling nucleus is detection signature

$$E_r^{\text{max}} \simeq \frac{2E_\nu^2}{M} \simeq 50 \text{ keV}$$

PHYSICAL REVIEW D

VOLUME 9, NUMBER 5

1 MARCH 1974



Coherent effects of a weak neutral current

Daniel Z. Freedman[†]

National Accelerator Laboratory, Batavia, Illinois 60510


and Institute for Theoretical Physics, State University of New York, Stony Brook, New York 11790

(Received 15 October 1973; revised manuscript received 19 November 1973)



CE ν NS Cross Section

- CE ν NS is the dominant cross section at low neutrino energy
- As neutrino energy increases, incoherent processes increase
- Total coherent cross section is linear in neutrino energy at low energy



Structure CE ν NS Signal

- Standard Model Prediction

$$\frac{d\sigma}{dE} = \frac{G_F^2}{4\pi} \left[(1 - 4 \sin^2 \theta_w) Z - N \right]^2 M \left(1 - \frac{ME}{2E_\nu^2} \right) F(Q^2)^2$$


$\approx 0 \rightarrow$ protons have small effect

square of sum \rightarrow part of coherence condition

nuclear form factor
 \rightarrow distribution of neutrons

- Recoil energy (M^{-1}) and rate (N^2)


Detection Rate [ton $^{-1}$ year $^{-1}$]¹



¹K. Scholberg, *Phys. Rev. D* **73** (2006) 033005. arXiv:hep-ex/0511042.

Physics Motivations for CE ν NS

- Non-Standard Interactions
- Multiple targets can greatly improve CHARM experiment limits on NSI
- Meaningful limits can be set with first generation experiments
- NSI may have significant influence in DUNE CP-violation search¹



$$\mathcal{L}_{\nu H}^{NSI} = -\frac{G_F}{\sqrt{2}} \sum_{\substack{q=u,d \\ \alpha,\beta=e,\mu,\tau}} [\bar{\nu}_\alpha \gamma^\mu (1 - \gamma^5) \nu_\beta] \times (\varepsilon_{\alpha\beta}^{qL} [\bar{q} \gamma_\mu (1 - \gamma^5) q] + \varepsilon_{\alpha\beta}^{qR} [\bar{q} \gamma_\mu (1 + \gamma^5) q])$$

(Plot) K. Scholberg, *Phys. Rev.* **D73** (2006) 033005. [arXiv:hep-ex/0511042](https://arxiv.org/abs/hep-ex/0511042).

¹M. Masud, A. Chatterjee, P. Mehta, [arXiv:1510.08261](https://arxiv.org/abs/1510.08261) [hep-ph].

Physics Motivations for CEνNS

- Non-Standard Interactions
- Multiple targets can greatly improve CHARM experiment limits on NSI
- Meaningful limits can be set with first generation experiments
- NSI may have significant influence in DUNE CP-violation search¹

TABLE I. Constraints on NSI parameters, from Ref. [35].

NSI parameter limit	Source
$-1 < \varepsilon_{ee}^{uL} < 0.3$	CHARM $\nu_e N$, $\bar{\nu}_e N$ scattering
$-0.4 < \varepsilon_{ee}^{uR} < 0.7$	
$-0.3 < \varepsilon_{ee}^{dL} < 0.3$	CHARM $\nu_e N$, $\bar{\nu}_e N$ scattering
$-0.6 < \varepsilon_{ee}^{dR} < 0.5$	
$ \varepsilon_{\mu\mu}^{uL} < 0.003$	NuTeV νN , $\bar{\nu} N$ scattering
$-0.008 < \varepsilon_{\mu\mu}^{uR} < 0.003$	
$ \varepsilon_{\mu\mu}^{dL} < 0.003$	NuTeV νN , $\bar{\nu} N$ scattering
$-0.008 < \varepsilon_{\mu\mu}^{dR} < 0.015$	
$ \varepsilon_{e\mu}^{uP} < 7.7 \times 10^{-4}$	$\mu \rightarrow e$ conversion on nuclei
$ \varepsilon_{e\mu}^{dP} < 7.7 \times 10^{-4}$	$\mu \rightarrow e$ conversion on nuclei
$ \varepsilon_{e\tau}^{uP} < 0.5$	CHARM $\nu_e N$, $\bar{\nu}_e N$ scattering
$ \varepsilon_{e\tau}^{dP} < 0.5$	CHARM $\nu_e N$, $\bar{\nu}_e N$ scattering
$ \varepsilon_{\mu\tau}^{uP} < 0.05$	NuTeV νN , $\bar{\nu} N$ scattering
$ \varepsilon_{\mu\tau}^{dP} < 0.05$	NuTeV νN , $\bar{\nu} N$ scattering


$$\mathcal{L}_{\nu H}^{NSI} = -\frac{G_F}{\sqrt{2}} \sum_{\substack{q=u,d \\ \alpha,\beta=e,\mu,\tau}} [\bar{\nu}_\alpha \gamma^\mu (1 - \gamma^5) \nu_\beta] \times [\varepsilon_{\alpha\beta}^{qL} \bar{q} \gamma_\mu (1 - \gamma^5) q] + [\varepsilon_{\alpha\beta}^{qR} \bar{q} \gamma_\mu (1 + \gamma^5) q]$$

(Plot) K. Scholberg, *Phys. Rev. D* **73** (2006) 033005. [arXiv:hep-ex/0511042](https://arxiv.org/abs/hep-ex/0511042).

¹M. Masud, A. Chatterjee, P. Mehta, [arXiv:1510.08261](https://arxiv.org/abs/1510.08261) [hep-ph].


Physics Motivations for CE ν NS

- Neutrino oscillations can use NC, flavor-blind CE ν NS interaction to look for L/E shape distortions and disappearance






(Left Plot) A.J. Anderson et al., *Phys. Rev.* **D86** (2012) 013004. arXiv:1201.3805 [hep-ph].

(Right Plot) J. Dutta et al., arXiv:1511.02834 [hep-ph].



Physics Motivations for CE ν NS

- CE ν NS is irreducible background for deep underground WIMP Dark Matter searches
- Supernova cross sections (or direct detection) via CE ν NS




(Left Plot) J. Billard, E. Figueroa-Feliciano, L. Strigari. arXiv:1307.5458 [hep-ph].

(Right Plot) K. Scholberg, *Phys. Rev. D* **73** (2006) 033005. arXiv:hep-ex/0511042.

¹A. Drukier and L. Stodolsky, *Phys. Rev. D* **30** (1984) 2295.



More Physics Motivations for CE ν NS

- Method to measure $\sin^2 \theta_W$
- Neutrino magnetic moment distorts recoil spectrum
- Low-mass Dark Matter searches
- Nuclear physics form factors
- Nuclear safeguards




(Bottom-Left Plot) K. Patton et al., *Phys. Rev.* **C86** (2012) 024612 arXiv:1207.0693 [nucl-th].

(Bottom-Right Plot) Horowitz et al, *Phys. Rev.* **D86** (2012) 013004. arXiv:1201.3805 [hep-ph].




Neutrino Production at the SNS

- 0.9-1.3 GeV protons on liquid mercury target produces π^+
- Total power: 0.9–1.4 MW





- Pulsed at 60 Hz for ~400 ns
→ $\text{few} \times 10^{-4}$ duty (steady-state background reduction)
- 43×10^6 v/cm²/s 20 m at SNS maximum power (1.4 MW)




Neutrino Production at the SNS

- 0.9-1.3 GeV protons on liquid mercury target produces π^+

- Total power: 0.9–1.4 MW





- Pulsed at 60 Hz for ~400 ns
→ $\text{few} \times 10^{-4}$ duty (steady-state background reduction)
- 43×10^6 $\nu/\text{cm}^2/\text{s}$ 20 m at SNS maximum power (1.4 MW)



Aside: Why Not a Reactor?


- High flux of electron antineutrinos is very alluring $\sim 10^{20}$
- Energy spectrum is few-MeV
- Cross section goes as E_ν
- Recoil energy goes as E_ν^2
- No pulsing, just reactor on/off cycle to measure backgrounds



¹H. Murayama & A. Pierce, *Phys. Rev. D* **65** (2002) 013012. [arXiv:hep-ph/0012075](https://arxiv.org/abs/hep-ph/0012075).


Aside: Why Not a Reactor?

- High flux of electron antineutrinos is very alluring $\sim 10^{20}$
- Energy spectrum is few-MeV
- Cross section goes as E_ν
- Recoil energy goes as E_ν^2
- No pulsing, just reactor on/off cycle to measure backgrounds



¹H. Murayama & A. Pierce, *Phys. Rev. D* **65** (2002) 013012. [arXiv:hep-ph/0012075](https://arxiv.org/abs/hep-ph/0012075).

Comparing “Stopped” Pion Sources




Higher energy accelerators begin producing kaons and this leads to “non-coherent” 236 MeV neutrinos

¹Plot from K. Scholberg

Physics Goals for COHERENT



- Measure N^2 dependence across multiple targets
- Deploy detectors in low-neutron-background basement area
- Measure relevant neutrino-induced neutrons and quenching factors

Nuclear Target	Technology	Mass (kg)	Distance from source (m)	Recoil threshold (keV _{nr})	Data-taking start date; CEvNS detection goal
CsI[Na]	Scintillating Crystal	14	20	6.5	9/2015; 3σ in 2 yr
Ge	HPGe PPC	10	22	5	Fall 2016
LAr	Single-phase scintillation	35	29	20	Fall 2016
NaI[Tl]	Scintillating crystal	185 [*] /2000	28	13	[*] high-threshold deployment started July 2016




Physics Goals for COHERENT

- Measure N^2 dependence across multiple targets
- Deploy detectors in low-neutron-background basement area
- Measure relevant neutrino-induced neutrons and quenching factors







Neutrino “Alley”

- Utility corridor has 22.5 m² for neutrino experiments and is largely decoupled from SNS operations; accessible via truck bay
- 1.5 m concrete floor supports large shielding structures
- 8 m.w.e. overburden and engineered backfill → neutron quiet
- 40 kW dedicated power install in October 2016



Detector Subsystems: CsI[Na]

- Na doping reduces afterglow seen in common TI doping
 - TI doping insufficient for near-surface experiment
- Cs and I surround Xe in periodic table, very similar nuclear recoil response
- Statistical nuclear-/electron-recoil separation
- Quenching factors measured
→ 5 keVnr is easily achieved





2 kg test crystal. Screened salts, electroformed Cu can (PNNL), ULB window and reflector.

¹J.I. Collar et al., *Nucl. Instrum. Meth. A* **773** (2015) 56.

Detector Subsystems: CsI[Na]


- 14 kg installed and running since July 2015 in neutrino alley
- Pb, water, and plastic shield
- Significant work on quenching factors has improved shows CE ν NS is within reach
- Steady state backgrounds are 10-20% less at ORNL
- Neutrino-induced neutron backgrounds reduced to 4% with improved HDPE shield
- Results soon!



¹J.I. Collar et al., *Nucl. Instrum. Meth. A* **773** (2015) 56.

Detector Subsystems: NaI[Tl]

- Born from discontinued DHS program; ~8 tons of NaI available
- 185 kg prototype for initial deployment
- 2 ton next phase deployment
- Up to 9 tons available
- NC $\text{CE}_{\nu\text{NS}}$ interaction
- Also CC interaction with ν_e
- https://twitter.com/NaIvE_SNS



multi-ton concept




^{55}Fe	$^{55}\text{Fe}(\nu_e, e^-)^{55}\text{Co}$	Stopped π/μ	KARMEN	$256 \pm 108(\text{stat}) \pm 43(\text{sys})$	264 [Shell] (Kolbe <i>et al.</i> , 1999a)
^{71}Ga	$^{71}\text{Ga}(\nu_e, e^-)^{71}\text{Ge}$	^{51}Cr source	GALLEX, ave.	$0.0054 \pm 0.0009(\text{tot})$	0.0058 [Shell] (Haxton, 1998)
		^{51}Cr	SAGE	$0.0055 \pm 0.0007(\text{tot})$	
		^{37}Ar source	SAGE	$0.0055 \pm 0.0006(\text{tot})$	0.0070 [Shell] (Bahcall, 1997)
^{127}I	$^{127}\text{I}(\nu_e, e^-)^{127}\text{Xe}$	Stopped π/μ	LSND	$284 \pm 91(\text{stat}) \pm 25(\text{sys})$	210-310 [Quasi-particle] (Engel <i>et al.</i> , 1994)

¹J.A. Formaggio and G. Zeller, *Rev. Mod. Phys.* **84** (2012) 1307.

Detector Subsystems: Liquid Argon

- Single-phase, scintillation only
- Built at FNAL by J. Yoo et al.
- 35-kg fiducial volume
- Readout is $2 \times$ Hamamatsu R5912-02MOD PMT (8" cryogenic, high-gain)
- Excellent nuclear-/electron-recoil PSD demonstrated by miniCLEAN
- SCENE has measured quenching factors¹
- ^{39}Ar helped by PSD and duty factor



¹H. Cao et al., SCENE Collaboration, *Phys. Rev. D* **91** (2015) 092007. [arXiv:1406.4825](https://arxiv.org/abs/1406.4825) [physics.ins-det].


Detector Subsystems: Liquid Argon

- Initial construction at Indiana University in Summer 2016
- Detector moved to SNS
- Construction is underway
- Running expected in December
- SNS run will contain shielding (inner layer to outer layer):
 - 23 cm water
 - 1/2" Cu
 - 4" Pb




Detector Subsystems: Liquid Argon

- Initial construction at Indiana University in Summer 2016
- Detector moved to SNS
- Construction is underway
- Running expected in December
- SNS run will contain shielding (inner layer to outer layer):
 - 23 cm water
 - 1/2" Cu
 - 4" Pb







Detector Subsystems: Liquid Argon

- Initial construction at Indiana University in Summer 2016
- Detector moved to SNS
- Construction is underway
- Running expected in December
- SNS run will contain shielding (inner layer to outer layer):
 - 23 cm water
 - 1/2" Cu
 - 4" Pb




Detector Subsystems: Germanium

- HPGe PPC type that have excellent low-energy resolution and low threshold
- Repurposing of Majorana Demonstrator and LANL ^{nat}Ge detectors
- First phase is 5-10 kg of existing BeGe detectors in a copper, lead, and polyethylene shielding system
- Second phase could add more detector mass with larger (C4-style) detectors
- Quenching factors well known



Neutron Backgrounds: Sandia NSC



- Sandia Neutron Scatter Camera
- 2 Planes of EJ-309 neutron detectors (neutron/gamma PSD)
- Reconstruct neutron direction and energy
- Optimizable plane spacing



¹N. Mascarenhas et al., *IEEE Trans. Nucl. Sci.* **56** (20109) 1269.




Neutron Backgrounds: Sandia NSC

- Mapped neutrons across SNS
- Significant rate drop from guide hall to basement locations
- “Prompt” component visible
- “Delayed” neutrons in basement significantly suppressed



Neutron Backgrounds: SciBath



- 80 L liquid scintillator
768 WLS fibers readout
→ event topology
- No optical separation
and uniform tracking
efficiency



¹R. Tayloe, *Nucl. Instrum. Meth.* **A562** (2006) 198.



Neutron Backgrounds: SciBath

- Measured at LAr site in SNS Neutrino Alley
- **Prompt flux (5-30 MeV) $\rightarrow (2.1 \pm 0.4) \times 10^{-5} \text{ m}^{-2} \mu\text{s}^{-1} \text{ MW}^{-1}$**
- Delayed flux $\rightarrow (1.9 \pm 0.7) \times 10^{-5} \text{ m}^{-2} \mu\text{s}^{-1} \text{ MW}^{-1}$
- Fluka transport neutrons $\rightarrow (3.2 \pm 0.3)$ in ROI per year (prompt)




Neutrino Induced Neutrons (NINs)

- Pb shielding interacts with ν_e and can break-up to 1 n or 2 n
- Shield can give CE ν NS backgrounds
- “Neutrino Cubes” testing neutron production from Pb, Fe inside water shield
- <https://twitter.com/theLeadNube>




Quenching Measurements

- Chicago has used commercial D-D neutron generator
- Duke group at TUNL have created a tunable $p\text{-}^7\text{Li}$ and D-D neutron sources to test low-energy nuclear recoils for quenching factor measurements






TUNL Setup



Chicago Setup

Conclusions

- COHERENT is deploying a suite of detector technologies in the neutron-quiet SNS basement
- First light measurement to test N^2 dependence
- Sensitivities shown below (LAr is being completed)
- LAr expects $\sim 300/700$ events on $\sim 100/900$ steady-state backgrounds in a year in prompt/delayed region



The COHERENT collaboration

arXiv:1509.08702



Institution	Board Member
University of California, Berkeley	Kai Vetter
University of Chicago	Juan Collar
Duke University	Kate Scholberg
University of Florida	Heather Ray
Indiana University	Rex Tayloe
Institute for Theoretical and Experimental Physics, Moscow	Dmitri Akimov
Lawrence Berkeley National Laboratory	Ren Cooper
Los Alamos National Laboratory	Steve Elliott
National Research Nuclear University MEPhI	Alex Bolozdynya
New Mexico State University	Robert Cooper
North Carolina Central University	Diane Markoff
North Carolina State University	Matt Green
Oak Ridge National Laboratory	Jason Newby
Sandia National Laboratories	David Reyna
University of Tennessee, Knoxville	Yuri Efremenko
Triangle Universities Nuclear Laboratory	Phil Barbeau
University of Washington	Jason Detwiler



- Collaboration: ~65 members, 16 institutions (USA+ Russia)

